



# Comparison of HIRDLS L1 and L2 Data for T, O<sub>3</sub> and H<sub>2</sub>O with ECMWF Analyses and Derived Radiances

Brian Kerridge, Alison Waterfall, Jolyon Reburn

Remote-Sensing Group
Earth Observation and Atmospheric Science Division, SSTD
Rutherford Appleton Laboratory, UK





#### Content



## Validation Activities at RAL

- Outline of Work
- Correlative Data
- Relevant Instrument Details
- Radiance Comparisons (L1)
- Product Comparisons (L2)
- Summary and Conclusion







#### **Outline**



- The work carried out at RAL covers two aspects:
  - Examination of HIRDLS corrected radiances, L1/HIRRAD, for channels designated for T, H<sub>2</sub>O and O<sub>3</sub>
    - Comparison with simulated radiances using ECMWF met analysis fields
  - Examination of HIRDLS L2 T and O<sub>3</sub>
    - Comparison to ECMWF geophysical data and sonde data





#### **Correlative Data**



#### **ECMWF** Analyses

## Temperature, H<sub>2</sub>O and O<sub>3</sub> data

- Taken on a regular 1.125° x 1.125° grid in lat/lon, on model levels (60 levels up to 0.1 hPa)
- Data available at 4 times throughout the day (00.00, 06.00, 12.00 and 18.00 UTC)
- For comparisons, data interpolated spatially and in time to the locations of HIRDLS measurements







#### **Correlative Data**



#### Sonde Data

#### Temperature profile data

- UKMO high resolution radiosondes obtained from the BADC
- 9 stations worldwide (many around UK)
- Matching criteria: 300 km & 3 hrs







#### Instrument



Temperature

Ozone

Water Vapour

Channel	Species	50% Response (cm <sup>-1</sup> )		Sounding Range	Radiometric Noise
		Lower	Upper	(km)	$(10^{-4}\mathrm{Wm^{-2}sr^{-1}})$
1	$N_2O, A$	$563.50 \pm 2.0$	$587.25 \pm 1.0$	8-70	12.0
2	CO <sub>2</sub> -L	$600.50 \pm 2.0$	$614.75 \pm 1.0$	8–40	6.3
3	CO <sub>2</sub> -M	$610.00 \pm 3.0$	$639.50 \pm 2.0$	8–60	5.9
4	CO <sub>2</sub> -M	$626.00 \pm 3.0$	$660.00 \pm 3.0$	15–60	6.0
5	CO <sub>2</sub> -H	$655.00 \pm 3.0$	$680.00 \pm 2.0$	30–105	4.3
6	A	$821.50 \pm 2.3$	$835.00 \pm 2.4$	8–55	1.9
7	CFC11	$835.00 \pm 2.4$	$852.00 \pm 2.4$	8-50	2.0
8	HNO <sub>3</sub>	$861.50 \pm 2.5$	$903.50 \pm 2.5$	8–70	4.2
9	CFC12	$916.00 \pm 2.6$	$931.50 \pm 2.6$	8-50	2.0
10	O <sub>3</sub> -M	$991.00 \pm 2.8$	$1009.00 \pm 2.8$	8–55	1.5
11	О3-Н	$1011.00 \pm 2.9$	$1046.50 \pm 2.9$	30–85	2.4
12	O <sub>3</sub> -L	$1120.00 \pm 3.2$	$1138.50 \pm 3.2$	8–55	0.96
13	A	$1202.00 \pm 3.4$	$1259.75 \pm 3.4$	8–55	1.1
14	$N_2O_5$	$1229.50 \pm 2.0$	$1259.75 \pm 1.0$	8–60	1.1
15	N <sub>2</sub> O	$1256.25 \pm 1.0$	$1281.75 \pm 1.0$	8-70	1.1
16	ClONO <sub>2</sub>	$1278.25 \pm 1.0$	$1298.75 \pm 1.0$	8-70	1.1
17	CH <sub>4</sub>	$1325.50 \pm 3.8$	$1367.50 \pm 3.8$	8–80	1.2
18	H <sub>2</sub> O-L	$1387.00 \pm 4.0$	$1435.00 \pm 4.0$	8-40	1.2
19	A	$1402.25 \pm 1.0$	$1415.75 \pm 1.0$	8–55	1.3
20	H <sub>2</sub> O-H	$1422.00 \pm 4.1$	$1542.00 \pm 4.3$	15–85	1.6
21	$NO_2$	$1585.50 \pm 4.5$	$1630.50 \pm 4.6$	8-70	1.1






#### Instrument



#### **HIRDLS Channel Filters**










#### Instrument



#### Field of View Map

**Space** 

Earth



4.50 mm 55.0 km 18.2 mrad







#### L1 Comparisons



#### Calculations

- Radiances generated using the HIRDLS FM for May 2006 period
- Channels
  - 2 5 (temperature)
  - 10 12 (ozone)
  - 18 & 20 (water vapour)
- Calculations based on the HIRRAD data files
- Temperature, H<sub>2</sub>O and O<sub>3</sub> taken from ECMWF, other contaminants from HIRDLS climatology files


Note: Cloud not included in the simulations and aerosol only as simple cilmatology so there will be tropospheric deviations of HIRDLS measurements for some channels





# Temperature Channel 2










## Temperature Channel 2








#### Ozone Channel 11










#### Ozone **Channel 11**












# Water Vapour Channel 20










# Water Vapour Channel 20







#### L2 Comparisons



#### **ECMWF** Data

 L2 data compared directly with ECMWF analysis data interpolated spatially and temporally to HIRDLS profile positions

#### **HIRDLS Data**

Data versions V2.00 & V2.02

#### Sonde Data

UKMO high resolution radiosondes






#### **Temperature**




#### Temperature - Global Average

Day Range : 20060504 - 20060531

(Shift: -1)














## Temperature



HIRDLS Temperature at 28.7299 hPa










#### Temperature












#### Ozone




#### O3 - Global Average

Day Range : 20060504 - 20060531

(Shift: -2)











#### **Conclusions**



#### **Summary & Conclusions**

- Radiance Comparisons
  - Radiance profiles have been simulated using the HIRDLS FM and ECMWF data
  - HIRDLS radiances show atmospheric structure predicted from the ECMWF data
- Product Comparisons
  - HIRDLS L2 T and O<sub>3</sub> results have been compared to data from sondes and ECMWF
  - Temperature data shows good agreement with correlative profiles; ~1-2K bias in the stratosphere
  - O<sub>3</sub> profiles agree at the 1ppmv level; ~1ppmv bias at peak

